

Practice		Population														Drug Problem																		
<p><i>Click on intervention title for a detailed description.</i></p> <p>Manual availability: <i>D = Download free from web</i> <i>F = Free print copy (order)</i> <i>\$ = Cost to purchase print copy</i> <i>N = No specific manual</i></p>	Manual: see note at left	Adolescents	Adults	African American	Am Indian/Alask Native	Asian/Pacific Islander	Children/Youth	College Student	Co-Occurring Patients	GLBT	Hispanic/Latino	HIV +/-Hep C/STD	Homeless	IV Drug User	Low-income/Unemployed	Men	Offenders	Opiate Subst Clients	Polydrug Users	Pregnant Women	Women	Not Specific to One Drug	Alcohol Dependence	Alcohol Problem Drinking	Amphetamines	Cocaine / Crack	Heroin / Opiates	Marijuana	Methamphetamine	Polydrug Use	Prescription Meds	Tobacco		
	Methadone Maintenance Treatment	N		X													X						X											
	Naltrexone (for Alcohol)	\$6		X																				X										
	Naltrexone (for Opiates)	N		X														X					X											

*Washington State law requires that a behavioral therapy component is provided to patients receiving pharmacological therapies for substance use disorders.

CITATION: University of Washington Alcohol and Drug Abuse Institute. Evidence-Based Practices for Treating Substance Use Disorders: Matrix of Interventions, August 2006.
 URL: <http://ada.i.washington.edu/ebp/matrix.pdf>