LOW-DOSE BUPRENORPHINE START

	DAY 1	DAY 2	DAY 3	DAY 4	DAY 5	DAY 6
	//	//	//	//	//	
How much do I take of a 2 mg strip?	1/4 strip daily	1/2 strip daily	1 strip daily	2 strips daily	3 strips daily	Come back to clinic for a check-in and refill.

What else can I take if I feel sick?

viiat eise caii i take ii i ieei sick:		
☐ clonidine every 8 hours as needed for withdrawal		
☐ diphenhydramine (Benadryl) every 6 hours as needed for withdrawal	Call	
☐ ondansetron (Zofran) every 8 hours as needed for nausea	if you have any	
☐ loperamide (Imodium) every 4 hours as needed for diarrhea		
☐ ibuprofen OR acetaminophen every 8 hours as needed for pain	questions	
☐ cyclobenzaprine every 8 hours as needed for spasms		
☐ trazodone at bedtime as needed for insomnia		

LOW-DOSE BUPRENORPHINE START

	DAY 6	DAY 7	DAY 8	DAY 9 +
	//	//	//	//
How much do I take of an 8 mg strip?	1 strip daily	1 & 1/2 strips daily	2 strips daily	Keep taking as instructed.

What else can I take if I feel sick?

☐ clonidine every 8 hours as needed for withdrawal	
☐ diphenhydramine (Benadryl) every 6 hours as needed for withdraw	/al
☐ ondansetron (Zofran) every 8 hours as needed for nausea	
☐ loperamide (Imodium) every 4 hours as needed for diarrhea	
☐ ibuprofen OR acetaminophen every 8 hours as needed for pain	
☐ cyclobenzaprine every 8 hours as needed for spasms	
☐ trazodone at bedtime as needed for insomnia	

Call (206) 744-2951 if you have any questions